Research Article

Non-adherence among women enrolled in a contraceptive vaginal ring use study in Kisumu, Kenya, 2014-2015

Eleanor McLellan-Lemal¹, Roman Gvetadze¹, Mitesh A Desai¹, Esther M Makanga², Yi Pan¹, Richard E Haaland¹, Angela N Holder¹, Victor Mudhune², Tiffany Williams¹, Taraz Samandari¹

¹ Centers for Disease Control and Prevention, Office of Infectious Diseases, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention, Division of HIV/AIDS Prevention, Atlanta, Georgia, USA, ² Kenya Medical Research Institute, Center for Global Health Research, Kisumu, Kenya

Keywords: global health

<https://doi.org/10.29392/joghr.2.e2018032>

Journal of Global Health Reports

Vol. 2, 2018

Background

Given future potential use of vaginal rings to prevent HIV infection, we examined the association of contraceptive vaginal ring (CVR) non-adherence with user dissatisfaction, tolerability, demographic, and behavioral factors.

Methods

In an open-label single-group study, sexually active women aged 18-34 years using oral or injectable hormonal contraception, conveniently sampled from general population, were assigned to 6-month use of a commercial CVR currently not licensed for use in Kenya. Non-adherence in any CVR cycle completed was assessed from: (1) self-report (not used for at least 1 day), and (2) pharmacy record (failure to timely receive a new CVR or return a used one). Additionally, non-adherence was assessed in a subset of participants by residual progestin and estrogen levels measured in returned CVRs.

Results

Of 202 participants who underwent CVR insertion by a study clinician, 142 completed all 6 visits, 172 responded to questions about ring use, and 43 provided used CVRs from months 1, 3, and 6 for residual hormone analysis. Non-adherence was 14.0% (24/172) by self-report and 54.5% (110/202) by pharmacy record. Non-adherence by pharmacy record was significantly reduced among women with a salary-based income (prevalence ratio (PR) 0.71, 95% confidence interval (CI) (0.55-0.91)] compared to women with income not salary-based or no income. Participants dissatisfied with CVR on ≥ 4 aspects (ambiguity of instructions, inconvenience of use, sensation, sexual discomfort, etc.) were more likely to report non-adherence (PR 2.69, 95% CI=(1.31-5.52)] compared to those dissatisfied with ≤ 3 aspects. Non-adherence by residual hormone levels was identified in 46.5% (20/43) participants. Over time, this subset of participants showed increasing non-adherence ($P=0.004$). We found lack of agreement among the various measures of non-adherence.

Conclusions

Economic empowerment interventions, especially those emphasizing partner-independent income options, and expanded education on CVR features may alleviate non-adherence. Addressing CVR dissatisfaction preemptively may also help mitigate non-adherence.

Vaginal rings that can deliver an antiretroviral (ARV) drug for preventing HIV infection, separately or in combination with agents for preventing pregnancy or sexually transmitted infections (STIs), are under development.¹⁻³ HIV prevention researchers anticipate that vaginal delivery of ARV drugs by long-acting (i.e., >28-day) methods will help avoid various adherence issues observed for some women with oral daily⁴ or peri-coitally dosed pre-exposure prophylaxis (PrEP).⁵ Contraceptive vaginal rings (CVRs) may offer valuable insights on users' experiences, including

potential adherence challenges associated with use of this technology, in particular among first-time users.

Worldwide, an overall low estimated prevalence (0.1%) of vaginal contraceptive methods, including CVRs, among women 15 to 49 years of age⁶ raises concerns about potential uptake (i.e., action taken to initiate use) and use of a multipurpose ring. Explanations for low uptake of CVRs include challenges with vaginal insertion, foreign-body sensation,⁷ sexual interference,⁸ and concerns about the ring getting lost inside the body.⁹ Poor CVR promotion, cost, and

short duration of use (e.g., 21-day use cycle) further account for CVR uptake obstacles.¹⁰ While high satisfaction, tolerability, favorable bleeding control, and adherence have been reported by women initiating CVR use,³ a continuation rate of 26% after 6 months by new CVR users has been observed, which is even lower than the 29% reported for combined oral contraceptives.¹¹

Adherence in clinical trials is subject to overestimation given challenges with measuring product use, both behaviorally and biologically.¹² Nonetheless, adherence in a trial is expected to be as good as or greater than “real world” use, where social, structural, and behavioral interventions may need to be coupled with biomedical ones and directed at providers and target populations alike.¹³ Women’s empowerment, defined as the “ability to make effective choices and to transform these choices into desired outcomes” as opposed to merely following prescriptive conventions,¹⁴ is a critical component in acceptability, satisfaction, and adherence of female-focused health interventions, including contraception. Women’s empowerment, which is influenced by individual, relational, and social determinants of health (e.g., socioeconomic status, education, gender dynamics, healthcare access, availability of methods, etc.), intersects with perceptions related to product properties, including the safety profile.

Given the seemingly apparent reasons that women would be interested in CVR use, it stands to reason that to optimize adherence, we need a better understanding of women’s experiences with this technology, in particular barriers to use. To address this issue, we examined the association of CVR non-adherence with user dissatisfaction, tolerability, demographic, and behavior variables among first-time users of this technology in Kisumu, Kenya. Of note, no CVR is licensed currently for use in Kenya, and the target population in this setting was predominantly naïve to CVRs. Our analysis sample included women who had at least one follow-up monthly visit post-CVR insertion. Two additional sub-analyses were performed for participants who provided follow-up behavioral data or were selected purposively for assessing residual progestin and estrogen levels in returned CVRs.

MATERIALS AND METHODS

Between April 2014 and August 2015, an open-label single-group study of NuvaRing® (Sharp & Dohme B.V., a subsidiary of Merck & Co., Inc, Kenilworth, NJ, USA) use was conducted in Kisumu, Kenya. In brief, a convenience sample of women was recruited from family planning and reproductive health clinics within Kisumu County via help of 10 community health volunteers and participant word-of-mouth referrals without incentives. We enrolled women who were 18-34 years of age, resided within a 150-kilometer radius of Kisumu City with no plans for relocation in the next 12 months, were fluent in English, Swahili, or Dholuo, provided documentation of depot medroxyprogesterone acetate (DMPA) or oral contraceptive pills (OCPs) use in the past three months, self-reported ≥ 2 episodes of vaginal intercourse on different days in the past 30 days at screening, tested negative for pregnancy and HIV, had no current or history of known medical contraindications for CVR use,

were not breastfeeding or within three months of parturition at screening, and were prepared to use the CVR for six months in place of injectable or oral contraceptives. Condom use for HIV and STI prevention was strongly encouraged.

Written informed consent was obtained prior to data and sample collection. Ring initiation schedules varied depending on completion of last OCP or DMPA-use cycle. OCP users were able to initiate ring use as soon as 30 days following study enrollment, while DMPA users may have had to delay ring initiation up to three months. To avoid overlap use in contraceptives, discontinuation of DMPA and initiation of NuvaRing® were recommended on the due date for the next injection. The informed consent process and data collection were available in the language choice of the participant (English, Swahili, or Dholuo). After receiving instruction and demonstration on a 3-dimensional female pelvic model, ring self-insertion and removal practice training occurred at the study clinic office for 210 women. At each follow-up visit, women received a bar of soap and 500 Kenya Shillings (approximately US\$ 5). In addition, as a part of the study clinic’s standard services, women received feminine sanitary pads, condoms, and hormonal contraceptives (at study exit or CVR discontinuation). Participants and their children were eligible to present to the study clinic at any time for diagnosis of common ailments and, if appropriate, referral for treatment. Participants’ sexual partners were entitled to receive free STI treatment following syndromic management assessment.

DATA COLLECTION

Non-adherence assessment was performed at follow-up visits scheduled between day 21 and day 29 of each one-month CVR cycle for six months following CVR initiation. At each visit, an electronic pharmacy log was used to record dates of each CVR dispensation and return of used CVRs by each participant. Of note, CVR dispensation and follow-up CVR visits occurred on the same date.

Demographic as well as baseline and quarterly behavioral data were collected using audio computer-assisted self-interview (ACASI). CVR user experiences and non-adherence were assessed using an adapted version of the NuvaRing® questionnaire developed by Novak et al,¹⁵ administered via computer-assisted personal interview (CAPI). Our questionnaire covered five broad CVR dimensions: difficulty of use, ambiguity of instructions, sexual discomfort, non-compliance, dissatisfaction. Willingness to recommend the CVR to others was also assessed. Within the dissatisfaction dimension, questions centered on specific CVR aspects (e.g., insertion, removal, placement, package use, physical comfort, partner support).

Testing for pregnancy was undertaken at each monthly visit. Pregnant women discontinued CVR use, received local antenatal care clinic referrals, and participated in quarterly follow-up. Rapid HIV testing was performed at baseline and every three months thereafter. Testing for other STIs and bacterial vaginosis was completed at baseline and study exit. Vaginal swabs were used to collect samples for gonorrhea, chlamydia, and bacterial vaginosis testing. Blood

samples were collected for herpes simplex virus type 2 (HSV-2) and syphilis testing.

MEASURES

Our primary outcomes were specified as binary variables summarizing different types of CVR non-adherence based on: (a) self-report, (b) pharmacy record, and (c) residual hormone levels. Participants were classified as non-adherent by self-report if they reported during follow-up any missed CVR use. Non-adherence by pharmacy record was determined if a new ring was not dispensed between days 19-31 for any CVR cycle or a used ring was not returned for every CVR cycle completed.

An objective measure of non-adherence was assessed by analyzing residual etonogestrel (progestin) and ethinyl estradiol (estrogen) levels in returned CVRs for months 1 ($n=26$), 3 ($n=43$), and 6 ($n=43$) for a subset of participants purposively selected from those who self-reported perfect (100%) adherence at all monthly visits (henceforward referred to as the hormone analysis sub-sample). NuvaRing® contains 11.7 milligrams (mg) of progestin, and 2.7 mg of estrogen. The average release rates per 24 hours over the 3-week use cycle for progestin and estrogen are 0.120 mg and 0.015 mg, respectively.¹⁶ Participants were classified as non-adherent if returned CVRs showed residual progestin or estrogen levels greater than 95% of those measured in a new, never used ring. Residual hormone levels were defined as consistent if the same use indicators were present across all returned CVRs examined (i.e., 100% overall use or 100% overall non-use).

Responses to demographic questions were coded as categorical variables. Incident STI or bacterial vaginosis, and responses to most risk behavior questions were included as binary variables. CVR dissatisfaction was characterized by a binary variable indicating displeasure with more than three CVR aspects within the four negative attitude domains, which included inconvenience of ring use (including package use), ambiguity of instructions, sexual discomfort, and difficulty with compliance. The number of dissatisfaction aspects reported at each CVR follow-up visit was used to visually depict dissatisfaction over time. Lastly, willingness to recommend the CVR to others was based a 5-point Likert agreement scale (highly agree, agree, undecided, disagree, and highly disagree).

Tolerability was summarized as a binary variable that distinguished between reports of ≥ 2 side effects (SEs) during the CVR-use period and reports of 1 or no SE. SEs included elevated systolic (>160 mmHg) or diastolic (>110 mmHg) blood pressure, self-reported fatigue, vaginal discharge, genital pain, headaches, depression, and abnormal vaginal bleeding. The number of SEs reported at each follow-up visit was also used to graphically depict tolerability over time.

STATISTICAL METHODS

We examined the factors associated with non-adherence among women who initiated ring use and completed at least one follow-up visit (non-adherence sample), in the sub-sample of participants who provided responses on ACASI

questionnaire (behavioral sub-sample), and in the sub-sample of participants included in the residual hormone-level analysis. Descriptive statistics were used to summarize categorical (frequency and percentage) and continuous (mean, median, standard deviation, and range) variables. The association with non-adherence outcomes was summarized by prevalence ratio (PR) and robust 95% confidence interval (CI) estimated from a log-binomial regression model using the generalized estimating equations (GEE) approach.^{17,18} The same GEE log-binomial regression with the visit number treated as a continuous covariate was used to model the over-time trend in a binary characteristic (dissatisfaction and tolerability). The incidence rate for pregnancy or HIV infection was estimated as the number of respective events occurred over study follow-up per 100 person-years with a robust 95% CI=obtained from a GEE Poisson model. Agreement among non-adherence measures was assessed by Cohen's kappa.¹⁹ All statistical tests were two-sided and interpreted at 0.05 level of significance. The analyses were performed in SAS 9.3 (SAS Institute Inc., Cary, NC, USA).

RESULTS

From the women pre-screened, 29.2% (202/692) initiated CVR use and completed at least one follow-up visit ([Figure 1](#)). Overall, three HIV seroconversions occurred during the study for an incidence rate of 3.6 per 100 person-years (95% CI=1.2-11.0) (data not shown). Among the non-adherence sample, 91.6% (185/202) who provided responses to ACASI questions were included in the behavioral sub-sample.

The non-adherence sample ($n=202$) accrued 83.7 out of the expected total 101 (82.9%) person-years of observation. Participants were followed for a median of 5.3 months, with follow-up ranging from 0.7 to 6.5 months (mean=5.0 months, standard deviation=1.1 months). Demographic characteristics of our CVR non-adherence sample are presented in [Table 1](#). Slightly over three-fourths of participants (79.7%) were using DMPA prior to initiating ring use.

Out of the 202 participants, 85.1% (172) self-reported on CVR non-adherence. Overall, 14.0% (24/172) and 54.5% (110/202) were non-adherent by self-report and by pharmacy record, respectively ([Table 1](#)). Over a total of 83.7 person-years of follow-up, five pregnancies occurred yielding a pregnancy rate of 6.0 per 100 person-years (95% CI=2.5-14.3) (data not shown).

The only significant factor associated with non-adherence by pharmacy record was the main source of personal income ([Table 1](#)). Specifically, the prevalence of non-adherence by pharmacy record was reduced by 29% among women with salary-based income compared to those with no personal income or no income (PR=0.71, 95% CI=0.55-0.91, $P=0.008$). CVR dissatisfaction was the only significant factor associated with non-adherence by self-report. Women indicating dissatisfaction with >3 CVR-related aspects (e.g., ring properties, use features) were more likely to be non-adherent by self-report compared to women dissatisfied with ≤ 3 aspects (PR=2.69 CI=1.31-5.52, $P=0.007$).

[Figure 2](#) depicts the proportion of participants ($n=202$) reporting any CVR dissatisfaction by study follow-up visit. Out of 491 dissatisfaction aspects reported across all follow-up visits, sexual discomfort accounted for 33.6% of dis-

Table 1. Self-reported and pharmacy record non-adherence by demographic, satisfaction, and tolerability factors (n=202), Kisumu contraceptive vaginal ring study, Kisumu, Kenya, 2014-2015*

Predictor	Non-adherence by self-report† n=24/172 (14.0%)					Non-adherence by pharmacy record‡ n=110/202 (54.5%)				
	Total, n (col %)	Non- adherent, n (col %)	Prevalence (%)	Prevalence ratio (95% CI)	P-value	Total, n (col %)	Non- adherent, n (col %)	Prevalence (%)	Prevalence ratio (95% CI)	P-value
Age at screening (years):					0.542					0.150
18-24	90 (52.3)	15 (62.5)	16.7	1.28 (0.40, 4.04)	0.677	104 (51.5)	52 (47.3)	50.0	1.04 (0.67, 1.61)	0.865
25-29	59 (34.3)	6 (25.0)	10.2	0.78 (0.21, 2.86)	0.707	71 (35.1)	45 (40.9)	63.4	1.32 (0.86, 2.02)	0.210
30-34	23 (13.4)	3 (12.5)	13.0	Ref.		27 (13.4)	13 (11.8)	48.1	Ref.	
Marital status:					0.459					0.559
Single	30 (17.9)	5 (21.7)	16.7	0.72 (0.20, 2.58)	0.617	31 (15.7)	19 (17.3)	61.3	1.43 (0.73, 2.78)	0.293
Married/cohabiting	125 (74.4)	15 (65.2)	12.0	0.52 (0.17, 1.56)	0.244	153 (77.3)	85 (77.3)	55.6	1.30 (0.70, 2.41)	0.413
Other	13 (7.7)	3 (13.0)	23.1	Ref.		14 (7.1)	6 (5.5)	42.9	Ref.	
Religion:					0.119					0.549
Roman Catholic	76 (44.4)	11 (47.8)	14.5	0.58 (0.24, 1.40)	0.225	89 (44.3)	51 (46.4)	57.3	0.97 (0.69, 1.35)	0.837
Other Christian	71 (41.5)	6 (26.1)	8.5	0.34 (0.12, 0.95)	0.040	80 (39.8)	40 (36.4)	50.0	0.84 (0.59, 1.21)	0.351
Other non-Christian	24 (14.0)	6 (26.1)	25.0	Ref.		32 (15.9)	19 (17.3)	59.4	Ref.	
Highest education completed:										
Primary or less	116 (68.2)	15 (65.2)	12.9	0.87 (0.39, 1.93)	0.737	135 (67.5)	76 (69.7)	56.3	1.11 (0.84, 1.47)	0.472
Secondary or higher	54 (31.8)	8 (34.8)	14.8	Ref.		65 (32.5)	33 (30.3)	50.8	Ref.	
Employment status:										
Employed	107 (62.9)	11 (47.8)	10.3	0.54 (0.25, 1.15)	0.110	126 (63.0)	66 (60.6)	52.4	0.90 (0.70, 1.16)	0.426
Unemployed	63 (37.1)	12 (52.2)	19.0	Ref.		74 (37.0)	43 (39.4)	58.1	Ref.	
Main source of personal income:										
Salary-based	88 (51.5)	9 (39.1)	10.2	0.61 (0.28, 1.33)	0.210	105 (52.2)	48 (43.6)	45.7	0.71 (0.55, 0.91)	0.008
Not salary-based/ None	83 (48.5)	14 (60.9)	16.9	Ref.		96 (47.8)	62 (56.4)	64.6	Ref.	
Parity:				N/A	N/A					0.407
0 live births	10 (5.9)	0 (0.0)	0.0			14 (7.0)	8 (7.3)	57.1	0.95 (0.58, 1.54)	0.831
1-2 live births	88 (51.8)	15 (65.2)	17.0			103 (51.5)	52 (47.3)	50.5	0.84 (0.65, 1.09)	0.181
3 or more live births	72 (42.4)	8 (34.8)	11.1			83 (41.5)	50 (45.5)	60.2	Ref.	

Non-adherence by self-report† n=24/172 (14.0%)						Non-adherence by pharmacy record‡ n=110/202 (54.5%)				
Hormonal contraceptive method at screening:										
DMPA	135 (78.5)	16 (66.7)	11.9	0.55 (0.25, 1.18)	0.124	161 (79.7)	87 (79.1)	54.0	0.96 (0.71, 1.31)	0.811
OCPs	37 (21.5)	8 (33.3)	21.6	Ref.		41 (20.3)	23 (20.9)	56.1	Ref.	
Age (years) of first sex:										
14 or less	35 (25.5)	7 (35.0)	20.0	1.57 (0.68, 3.62)	0.290	45 (27.8)	28 (32.6)	62.2	1.26 (0.94, 1.68)	0.127
15 or older	102 (74.5)	13 (65.0)	12.7	Ref.		117 (72.2)	58 (67.4)	49.6	Ref.	
History of unwanted/forced sex:										
Yes	57 (33.3)	9 (39.1)	15.8	1.29 (0.59, 2.79)	0.525	64 (31.8)	38 (34.5)	59.4	1.13 (0.87, 1.46)	0.353
No	114 (66.7)	14 (60.9)	12.3	Ref.		137 (68.2)	72 (65.5)	52.6	Ref.	
Pregnancy intentions in next 12 months:										
Intends/desires to get pregnant	8 (4.8)	1 (5.0)	12.5	1.05 (0.16, 6.91)	0.957	8 (4.7)	4 (5.2)	50.0	1.10 (0.54, 2.25)	0.788
No intent/desire to get pregnant	160 (95.2)	19 (95.0)	11.9	Ref.		161 (95.3)	73 (94.8)	45.3	Ref.	
Incident pregnancy:										
Yes	5 (2.9)	2 (8.3)	40.0	3.04 (0.97, 9.51)	0.057	5 (2.5)	2 (1.8)	40.0	0.73 (0.25, 2.15)	0.568
No	167 (97.1)	22 (91.7)	13.2	Ref.		197 (97.5)	108 (98.2)	54.8	Ref.	
Would recommend CVR to others:										
Agrees	154 (90.6)	23 (95.8)	14.9	2.39 (0.35, 16.54)	0.378	182 (91.9)	101 (93.5)	55.5	1.27 (0.72, 2.24)	0.414
Strongly agrees	16 (9.4)	1 (4.2)	6.3	Ref.		16 (8.1)	7 (6.5)	43.8	Ref.	
CVR dissatisfaction:§										
More than 3 dissatisfaction aspects	43 (30.5)	13 (54.2)	30.2	2.69 (1.31, 5.52)	0.007	47 (29.0)	27 (30.0)	57.4	1.05 (0.78, 1.41)	0.754
3 or less dissatisfaction aspects	98 (69.5)	11 (45.8)	11.2	Ref.		115 (71.0)	63 (70.0)	54.8	Ref.	
CVR tolerability:										
Side effect (SE) reported more than once	28 (16.8)	4 (18.2)	14.3	1.10 (0.40, 3.01)	0.848	34 (18.8)	18 (19.8)	52.9	1.07 (0.75, 1.52)	0.725
0 to 1 SE reported	139 (83.2)	18 (81.8)	12.9	Ref.		147 (81.2)	73 (80.2)	49.7	Ref.	
Attitude toward pelvic exam:										

Non-adherence by self-report† n=24/172 (14.0%)						Non-adherence by pharmacy record‡ n=110/202 (54.5%)				
Negative	26 (21.7)	4 (25.0)	15.4	1.21 (0.42, 3.43)	0.726	29 (20.4)	15 (19.2)	51.7	0.93 (0.63, 1.37)	0.705
Neutral or positive	94 (78.3)	12 (75.0)	12.8	Ref.		113 (79.6)	63 (80.8)	55.8	Ref.	

CI – confidence interval, DMPA - depot medroxyprogesterone acetate, OCPs – oral contraceptive pills, CVR – contraceptive vaginal ring

*Sample sizes fluctuate slightly for some variables due to missing data. Some percentages do not sum to 100 because of rounding.

†Due to incomplete or missing data, self-reported data was available for 85.1% (172/202) of the 202 CVR non-adherence sample.

‡Binary variable: a) new ring dispensation did not occur between days 19-31 for each cycle of use or (b) participant did not return a used ring for every CVR cycle completed. Four participants were solely classified as non-adherent by pharmacy record because they failed to return a used CVR for at least one follow-up visit.

§Dissatisfaction was assessed on 4 dimensions: inconvenience of ring use (including package use), ambiguity of instructions, sexual discomfort, and difficulty with compliance. Within each of these dimensions, several experiential or attitudinal questionnaire items (ie, aspects) were addressed.

Figure 1. Flowchart: Screening, enrollment, and follow-up, Kisumu contraceptive vaginal ring study, Kisumu, Kenya, 2014-2015.

*Information on participants who went from pre-screening to enrollment has been published elsewhere.¹⁵ †An additional participant was identified after study close out and preparation of final dataset. The number eligible for enrollment and the number enrolled differ by one from previously reported data,¹⁵ 304 and 302, respectively. ‡Among the 93 enrollees who did not initiate contraceptive vaginal ring (CVR) use, 43.0% withdrew their study participation, 45.2% were lost to follow-up, and 12.0% were no longer medically eligible. §Women initiating CVR use who completed at least one adherence follow-up visit; 8/210 (3.8%) did not have any follow-up time; 5 withdrew participation, 2 were medically discontinued, and 1 was lost to follow-up. ||Overall, 112 returned CVRs were assessed among 43 participants who self-reported perfect (100%) adherence at all monthly visits. At month 1, 26 returned CVRs were used to measure residual drug levels and to assess non-adherence.¹⁸ Forty-three CVRs were assessed at months 3 and 6. ¶CVR users who completed one or more ACASI questionnaires in which ≥1 month CVR experience occurred.

satisfactions (some participants may have reported this aspect at multiple visits).

SEs were reported by 9-14% of participants each month (Figure 3). The three most commonly reported SEs were headache, fatigue, and vaginal discharge. We found no significant associations between CVR tolerability and non-adherence either by self-report or by pharmacy record. All

reported SEs were mild in severity (Grade 1) and did not require treatment.

Table 2 displays non-adherence by self-report and by pharmacy record in the behavioral sub-sample (n=185). Out of the 185 participants, 84.3% (156) self-reported on CVR non-adherence. Relevant ACASI data were available for 19 of 24 participants who were non-adherent by self-report and 96.4% (106/110) of participants who were non-adherent

Figure 2. Proportion of participants (n=202) reporting ≥ 1 dissatisfaction aspect by Kisumu contraceptive vaginal ring study visit, Kisumu, Kenya, 2014-2015.

*Cumulative count of dissatisfaction aspects reported by all participants over study follow-up; the per participant mean represents the average cumulative count of dissatisfaction aspects reported over follow-up. †Other category includes lack of partner support (n=8), unclear instructions (n=5), and miscellaneous responses (n=25).

by pharmacy record. We found no evidence of an association between the behavioral measures and non-adherence by self-report or by pharmacy record (Table 2).

In the analysis of residual progestin and estrogen, a total of 112 returned CVRs were examined for 43 participants who self-reported perfect adherence at all monthly visits. Overall, 20 (46.5%) out of 43 participants were non-adherent by residual hormone levels, including 7 (16.3%) who showed consistent non-use and 13 (30.2%) who showed partial or inconclusive use (consistent with tampering). Analysis of residual hormone levels in returned CVRs showed increasing non-adherence over time ($P=0.004$, non-adherence for months 1, 3, and 6 were 19.2%, 32.6%, and 44.2%, respectively).

Finally, we observed poor to no agreement (Cohen's kappa = 0.03) between pharmacy record and self-reported non-adherence measures. There was disagreement observed between pharmacy record non-adherence and residual drug analysis as well as between self-reported non-adherence and residual drug analysis (data not shown).

DISCUSSION

We sought to understand the drivers of non-adherence to a CVR in a naïve population, with an intent to incorporate lessons learned into future studies of multipurpose HIV prevention rings in western Kenya. Despite all first-time users of CVR in our study being willing to recommend the ring to other women, which suggests that this technology for at least contraceptive purposes may be appropriate in this setting, following the guidelines for CVR use may have been a challenge for some, in particular those lacking a salary-based income or those being dissatisfied with >3 aspects of CVR use. Our sub-analysis of residual hormone levels in returned CVRs suggested increasing ring non-adherence over time. Among our behavioral sub-sample, no statistically significant factors for non-adherence either by self-report or by pharmacy record were found.

Clinical trials, observational studies, and medical practice have shown consistently high levels of satisfaction with NuvaRing®.^{20,21} In general, women have indicated the ring is easy to use, effective, and convenient, with interference with sexual intercourse and local SEs (e.g., leukorrhea, vaginal discomfort, vaginitis) as the primary reasons cited for

Table 2. Self-reported and pharmacy record non-adherence by sexually transmitted infections (STIs) and behavioral characteristics (n=185), Kisumu contraceptive vaginal ring study, Kisumu, Kenya, 2014-2015*

Predictor	Non-adherence by self-report n=19/156 (12.1%)					Non-adherence by pharmacy record† n=106/185 (57.3%)				
	Total, n (col %)	Non- adherent, n (%)	Prevalence (%)	Prevalence ratio (95% CI)	P-value	Total, n (col %)	Non- adherent, n (col %)	Prevalence (%)	Prevalence ratio (95% CI)	P-value
HIV incident infection:‡				N/A	N/A					
Yes	2 (1.3)	0 (0.0)	0.0			3 (1.6)	1 (0.9)	33.3	0.58 (0.12, 2.88)	0.503
No	154 (98.7)	19 (100.0)	12.3			182 (98.4)	105 (99.1)	57.7	Ref.	
Bacterial vaginosis incident infection:§				N/A	N/A					
Yes	13 (8.3)	0 (0.0)	0.0			14 (7.6)	7 (6.6)	50.0	0.86 (0.50, 1.48)	0.594
No	143 (91.7)	19 (100.0)	13.3			171 (92.4)	99 (93.4)	57.9	Ref.	
Herpes simplex virus type-2 (HSV-2) incident infection:§				N/A	N/A					
Yes	4 (2.6)	0 (0.0)	0.0			5 (2.7)	2 (1.9)	40.0	0.69 (0.23, 2.04)	0.505
No	152 (97.4)	19 (100.0)	12.5			180 (97.3)	104 (98.1)	57.8	Ref.	
Chlamydia incident infection§										
Yes	6 (3.8)	2 (10.5)	33.3	2.94 (0.87, 9.93)	0.082	6 (3.2)	3 (2.8)	50.0	0.87 (0.39, 1.95)	0.734
No	150 (96.2)	17 (89.5)	11.3	Ref.		179 (96.8)	103 (97.2)	57.5	Ref.	
Gonorrhea incident infection:§				N/A	N/A				N/A	N/A
Yes	1 (0.6)	0 (0.0)	0.0			2 (1.1)	2 (1.9)	100.0		
No	155 (99.4)	19 (100.0)	12.3			183 (98.9)	104 (98.1)	56.8		
Syphilis incident infection:§				N/A	N/A				N/A	N/A
Yes	0 (0.0)	0 (0.0)	0.0			0 (0.0)	0 (0.0)	0.0		
No	156 (100.0)	19 (100.0)	12.2			185 (100.0)	106 (100.0)	57.3		
Any STI or bacterial vaginosis incident infection:§										
Yes	24 (15.4)	2 (10.5)	8.3	0.65 (0.16, 2.62)	0.542	28 (15.1)	13 (12.3)	46.4	0.78 (0.52, 1.19)	0.254
No	132 (84.6)	17 (89.5)	12.9	Ref.		157 (84.9)	93 (87.7)	59.2	Ref.	
Number of sexual partners in past 3 months:					0.555					0.775
No partners	37 (24.3)	5 (26.3)	13.5	0.71 (0.21, 2.36)	0.575	44 (24.6)	26 (25.2)	59.1	1.18 (0.73, 1.92)	0.499
1 partner	94 (61.8)	10 (52.6)	10.6	0.56 (0.19, 1.61)	0.281	113 (63.1)	66 (64.1)	58.4	1.17 (0.75, 1.82)	0.495
2 or more partners	21 (13.8)	4 (21.1)	19.0	Ref.		22 (12.3)	11 (10.7)	50.0	Ref.	

Predictor	Non-adherence by self-report n=19/156 (12.1%)					Non-adherence by pharmacy record† n=106/185 (57.3%)				
	Total, n (col %)	Non- adherent, n (%)	Prevalence (%)	Prevalence ratio (95% CI)	P-value	Total, n (col %)	Non- adherent, n (col %)	Prevalence (%)	Prevalence ratio (95% CI)	P-value
Vaginal sex in past 3 months:										
Yes	117 (76.0)	14 (73.7)	12.0	0.89 (0.34, 2.29)	0.802	137 (75.7)	78 (75.0)	56.9	0.96 (0.72, 1.28)	0.799
No	37 (24.0)	5 (26.3)	13.5	Ref.		44 (24.3)	26 (25.0)	59.1	Ref.	
Vaginal sex without a condom in past 3 months:										
Yes	112 (72.7)	12 (63.2)	10.7	0.64 (0.27, 1.52)	0.315	130 (71.8)	72 (69.2)	55.4	0.88 (0.68, 1.15)	0.350
No	42 (27.3)	7 (36.8)	16.7	Ref.		51 (28.2)	32 (30.8)	62.7	Ref.	
Anal sex in past 3 months:										
Yes	55 (35.9)	4 (21.1)	7.3	0.48 (0.17, 1.36)	0.166	64 (35.6)	36 (34.6)	56.3	0.96 (0.74, 1.25)	0.760
No	98 (64.1)	15 (78.9)	15.3	Ref.		116 (64.4)	68 (65.4)	58.6	Ref.	
Anal sex without a condom in past 3 months:										
Yes	52 (34.2)	3 (15.8)	5.8	0.36 (0.11, 1.18)	0.092	59 (33.1)	33 (32.0)	55.9	0.95 (0.72, 1.25)	0.716
No	100 (65.8)	16 (84.2)	16.0	Ref.		119 (66.9)	70 (68.0)	58.8	Ref.	
HIV(+/unknown) partners in past 3 months:										
Yes	50 (33.6)	5 (26.3)	10.0	0.71 (0.27, 1.85)	0.481	57 (32.4)	28 (27.2)	49.1	0.78 (0.58, 1.05)	0.101
No	99 (66.4)	14 (73.7)	14.1	Ref.		119 (67.6)	75 (72.8)	63.0	Ref.	
Unwanted/forced sex in past 3 months:										
				N/A	N/A				N/A	N/A
Yes	6 (3.8)	0 (0.0)	0.0			7 (3.8)	7 (6.6)	100.0		
No	150 (96.2)	19 (100.0)	12.7			178 (96.2)	99 (93.4)	55.6		
Transactional sex or exchange sex in past 3 months:										
Yes	8 (5.1)	1 (5.3)	12.5	1.03 (0.16, 6.76)	0.977	9 (4.9)	3 (2.8)	33.3	0.57 (0.22, 1.45)	0.237
No	148 (94.9)	18 (94.7)	12.2	Ref.		176 (95.1)	103 (97.2)	58.5	Ref.	

CI – confidence interval, STI – sexually transmitted infection

*Behavioral sub-sample of participants who had completed one or more audio computer-assisted self-interview (ACASI) questionnaires and had used the contraceptive vaginal ring (CVR) for at least one month (n=185); relevant ACASI data were available for 19 out of the 24 participants non-adherent by self-report and 106 out of the 110 participants non-adherent by pharmacy record; sample sizes fluctuate slightly for some variables due to missing data. Some percentages do not sum to 100 because of rounding.

†Binary variable: a) new ring dispensation did not occur between days 19-31 for each cycle of use or (b) participant did not return a used ring for every CVR cycle completed. Four participants were solely classified as non-adherent by pharmacy record because they failed to return a used CVR for at least one follow-up visit.

‡Positive test at any time following CVR initiation.

§Laboratory-confirmed diagnoses at study exit for women completing at least one 28-day CVR cycle.

Figure 3. Proportion of participants (n=202) reporting ≥ 1 side effect (SE) by Kisumu contraceptive vaginal ring study visit, Kisumu, Kenya, 2014-2015.

*Cumulative count of SEs reported by all participants over study follow-up; the per participant mean represents the average cumulative count of SEs reported by a participant over follow-up. †Other category includes genital pain (n=4), depression (n=3), and elevated blood pressure (n=1).

either discontinuing or disliking the ring.^{21,22} Common reasons for disliking the CVR were similar for women in our trial. Presumably, with increased consumer education, as well as an improved understanding of user preferences, these barriers can be overcome.

Non-adherence ranging from 9% to 20% by self-report have been found in other NuvaRing® studies^{21,23} and was slightly lower (8%) in a recent dapivirine ring HIV prevention trial conducted in multiple African countries.²⁴ Our by-pharmacy-record measure may have under-estimated non-adherence. Other studies have defined NuvaRing® cycle non-adherence as a ring cycle that (a) extended 48 hours beyond day 22, and (b) the length of ring-free period was lengthened by more than 24 hours from day 8.²⁵ Notably, these studies collected diaries and tracking logs completed by the participant as opposed to our method, which involved calculations based on pharmacy dates for dispensation of new CVRs and return dates of a used CVRs. As suggested by others, non-adherence for longer-acting ARV rings require cumulative measurement over time that could benefit from non-invasive electronic and biometric monitoring technologies requiring minimal participant effort.²⁶

Other studies have found decreased non-adherence among women with an independent income.⁴ Women with

their own source of income may have greater family planning decision-making input than women who depend on their partners, or women who have variable income (eg, seasonal, casual, or temporary work). While additional research is needed, economic empowerment interventions suggest that most programs show reproductive health improvements.²⁷

A recent systematic review of contraceptive use in sub-Saharan Africa suggests that misinformation and concerns about perceived SEs are common barriers in accepting and using modern contraceptive methods.²⁸ As measured in our study, we did not find that SE concerns and tolerability were barriers to CVR use.

A number of limitations are associated with this study. The generalizability of our findings is limited due to our study design and sampling approach. While women in our sample were able to use a novel intravaginal ring to prevent pregnancy, which may have implications for future use of rings to prevent HIV and other STIs in this high-burden region, caution is warranted with such an interpretation.

A gold standard for assessing adherence to modern contraceptive methods is not available.²⁹ Our definitions of particular measures, in particular non-adherence by self-report, non-adherence by pharmacy record, may have pro-

duced imprecisions with either observation or measurement process. Measures for assessing CVR non-adherence in the literature are largely limited to self-report and subject to under-reporting of non-adherent behavior. We acknowledge that some women in our study could have had on-time CVR dispensation (between days 19 and 31 for each monthly cycle) and returned a used ring; yet, they could have been partially or completely non-adherent. Similar to self-report, a by-pharmacy-record measure is suboptimal in reliably assessing non-adherence. HIV risk perceptions along with HIV stigma, which were not measured in our study, may have created reticence to initiate ring use and influenced non-adherence behavior. Self-reported non-adherence and behavioral data may be subject to recall or social desirability biases. Performing residual hormone analysis on the returned rings from all women in our sample was not feasible. Moreover, such objective measures may not be entirely clear-cut in their ability to distinguish inconsistent users from consistent users.²⁹ Given small count, our pregnancy rate was estimated with low precision and reliability, and should be interpreted with caution. Lastly, NuvaRing® may offer a slight allowance for imperfect use given that removal of the ring for up to three hours for sexual or hygiene purposes does not compromise contraceptive efficacy.

CONCLUSIONS

While vaginal rings may be a highly effective but underutilized contraceptive approach and potentially a preferred future HIV prevention method for some women, perfect use will be a challenge. Apart from measurement challenges of non-adherence in clinical trial settings, real world non-adherence must be carefully accounted for in development of any multipurpose vaginal ring product. Economic empowerment interventions, in particular those that emphasize consistent and partner-independent income options, may mitigate non-adherence to modern contraceptive methods such as a CVR. To better understand non-adherent behavior, it may be necessary to go beyond looking for demographic and behavioral factors to explain CVR non-adherence, such as analyzing participants' relationship with power dynamics, self-assertion, and sexuality. Lastly, to further minimize non-adherence, preemptively addressing CVR dissatisfaction and expanding education on CVR features may be beneficial.

ACKNOWLEDGEMENTS

The authors would like to thank the study participants for their time and contribution to this study as well as the broader Kisumu community for its support. The authors would also like to thank Beatrice Nyagol, Kenneth Ondeng'e, Lawrence Opado, Richard Ndivo, and members of the study implementation team. The authors extend a special thanks to Siobhan O'Connor, Clyde Hart, and David Schnabel for their technical guidance and support.

DISCLAIMER

The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the United States Centers for Disease Control and Prevention (CDC) or the Department of Health and Human Services. The Kenya Medical Research Institute (KEMRI) Director and the KEMRI Publication Review Committee gave permission to publish this manuscript.

FUNDING

The Centers for Disease Control and Prevention supported this work under PN1757, 2014-2015.

CLINICAL TRIAL INFORMATION

The study is registered at [ClinicalTrials.gov](https://clinicaltrials.gov) under the identifier NCT02529683.

COMPETING INTERESTS

The authors have completed the Unified Competing Interest form at http://www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare no conflict of interest.

CORRESPONDENCE TO:

Eleanor McLellan-Lemal
Centers for Disease Control and Prevention
1600 Clifton Rd., MS-E45
Atlanta, GA 30329
United States of America
egm4@cdc.gov

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-4.0). View this license's legal deed at <http://creativecommons.org/licenses/by/4.0> and legal code at <http://creativecommons.org/licenses/by/4.0/legalcode> for more information.

REFERENCES

1. Sitruk-Ware R, Nath A, Mishell DR Jr. Contraception technology: past, present and future. *Contraception*. 2013;87(3):319-330. doi:10.1016/j.contraception.2012.08.002
2. Abdool Karim Q, Abdool Karim SS, Frohlich JA, et al. Effectiveness and safety of tenofovir gel, an antiretroviral microbicide, for the prevention of HIV infection in women. *Science*. 2010;329(5996):1168-1174. doi:10.1126/science.1193748
3. Bahamondes L, Bahamondes MV. New and emerging contraceptives: a state-of-the-art review. *Int J Womens Health*. 2014;6:221-234. doi:10.2147/ijwh.s46811
4. Marrazzo JM, Ramjee G, Richardson BA, et al. Tenofovir-based preexposure prophylaxis for HIV infection among African women. *N Engl J Med*. 2015;372(6):509-518. doi:10.1056/nejmoa1402269
5. Thurman A, Clark M, Hurlburt J, Doncel G. Intravaginal rings as delivery systems for microbicides and multipurpose prevention technologies. *Int J Womens Health*. 2013;5:695-708. doi:10.2147/ijwh.s34030
6. Guttmacher Institute. Contraceptive use in the United States fact sheet. Published 2016. Accessed April 30, 2018. <https://www.guttmacher.org/fact-sheet/contraceptive-use-united-states>
7. Oddsson K, Leifels-Fischer B, de Melo NR, et al. Efficacy and safety of a contraceptive vaginal ring (NuvaRing) compared with a combined oral contraceptive: a 1-year randomized trial. *Contraception*. 2005;71(3):176-182. doi:10.1016/j.contraception.2004.09.001
8. Stewart FH, Brown BA, Raine TR, Weitz TA, Harper CC. Adolescent and young women's experience with the vaginal ring and oral contraceptive pills. *J Pediatr Adolesc Gynecol*. 2007;20(6):345-351. doi:10.1016/j.jpag.2007.06.001
9. van der Straten A, Montgomery ET, Cheng H, et al. High acceptability of a vaginal ring intended as a microbicide delivery method for HIV prevention in African women. *AIDS Behav*. 2012;16(7):1775-1786. doi:10.1007/s10461-012-0215-0
10. Brache V, Payán LJ, Faundes A. Current status of contraceptive vaginal rings. *Contraception*. 2013;87(3):264-272. doi:10.1016/j.contraception.2012.08.037
11. Ahrendt HJ, Nisand I, Bastianelli C, et al. Efficacy, acceptability and tolerability of the combined contraceptive ring, NuvaRing, compared with an oral contraceptive containing 30 µg of ethinyl estradiol and 3 mg of drospirenone. *Contraception*. 2006;74(6):451-457. doi:10.1016/j.contraception.2006.07.004
12. Woodsong C, MacQueen K, Amico KR, et al. Microbicide clinical trial adherence: insights for introduction. *J Int AIDS Soc*. 2013;16:18505. doi:10.7448/ias.16.1.18505
13. Tolley EE, Severy LJ. Integrating behavioral and social science research into microbicide clinical trials: challenges and opportunities. *Am J Public Health*. 2006;96:79-83. doi:10.2105/ajph.2004.043471
14. World Bank. Promoting women's agency. In: *World Development Report 2012: Gender Equality and Development*. World Bank; 2012.
15. Novák A, de la Loge C, Abetz L. Development and validation of an acceptability and satisfaction questionnaire for a contraceptive vaginal ring, NuvaRing. *Pharmacoeconomics*. 2004;22(4):245-256. doi:10.2165/00019053-200422040-00003
16. Wieder DR, Pattimaki L. Examining the efficacy, safety, and patient acceptability of the combined contraceptive vaginal ring (NuvaRing). *Int J Womens Health*. 2010;2:401-409. doi:10.2147/ijwh.s6162
17. McNutt LA, Wu C, Xue X, Hafner JP. Estimating the relative risk in cohort studies and clinical trials of common outcomes. *Am J Epidemiol*. 2003;157(10):940-943. doi:10.1093/aje/kwg074
18. Hilbe JM, Hardin J, Hardin H. *Generalized Estimating Equations*. 2nd ed. CRC Press; 2003.
19. Cohen J. A coefficient of agreement for nominal scales. *Educ Psychol Meas*. 1960;20:37-46. doi:10.1177/001316446002000104
20. Roumen FJ, op ten Berg MM, Hoomans EH. The combined contraceptive vaginal ring (NuvaRing®): first experience in daily clinical practice in the Netherlands. *Eur J Contracept Reprod Health Care*. 2006;11:14-22. doi:10.1080/13625180500389547
21. Roumen FJ, Mishell DR Jr. The contraceptive vaginal ring, NuvaRing®, a decade after its introduction. *Eur J Contracept Reprod Health Care*. 2012;17(6):415-427. doi:10.3109/13625187.2012.713535

22. Roumen FJ. Review of the combined contraceptive vaginal ring, NuvaRing®. *Ther Clin Risk Manag.* 2008;4:441-451. [doi:10.2147/tcrm.s1964](https://doi.org/10.2147/tcrm.s1964)
23. Dieben TO, Roumen FJ, Apter D. Efficacy, cycle control, and user acceptability of a novel combined contraceptive vaginal ring. *Obstet Gynecol.* 2002;100:585-593.
24. Nel A, Bekker LG, Bukusi E, et al. Safety, acceptability and adherence of dapivirine vaginal ring in a microbicide clinical trial conducted in multiple countries in sub-Saharan Africa. *PLoS ONE.* 2016;11(3):e0147743. [doi:10.1371/journal.pone.0147743](https://doi.org/10.1371/journal.pone.0147743)
25. Bruni V, Pontello V, Luisi S, Petraglia F. An open-label, multicentre trial to evaluate the vaginal bleeding pattern of the combined contraceptive vaginal ring NuvaRing®. *Eur J Obstet Gynecol Reprod Biol.* 2008;139:65-71. [doi:10.1016/j.ejogrb.2008.02.001](https://doi.org/10.1016/j.ejogrb.2008.02.001)
26. MacQueen KM, Tolley EE, Owen DH, et al. An interdisciplinary framework for measuring and supporting adherence in HIV prevention trials of ARV-based vaginal rings. *J Int AIDS Soc.* 2014;17(3(Suppl 2)):19158. [doi:10.7448/ias.17.3.19158](https://doi.org/10.7448/ias.17.3.19158)
27. Reed E, West BS, Salazar M, Monroy RV. Economic empowerment to improve sexual and reproductive health among women and girls. In: *Global Perspectives on Women's Sexual and Reproductive Health across the Lifecourse.* Springer International Publishing; 2018:297-315. [doi:10.1007/978-3-319-60417-6_17](https://doi.org/10.1007/978-3-319-60417-6_17)
28. Blackstone SR, Nwaozuru U, Iwelunmor J. Factors influencing contraceptive use in sub-Saharan Africa: a systematic review. *Int Q Community Health Educ.* 2017;37(2):79-91. [doi:10.1177/0272684x16685254](https://doi.org/10.1177/0272684x16685254)
29. Smith C, Edwards P, Free C. Assessing the validity and reliability of self-report data on contraception use in the Mobile Technology for Improved Family Planning (MOTIF) randomised controlled trial. *Reprod Health.* 2018;15:50. [doi:10.1186/s12978-018-0494-7](https://doi.org/10.1186/s12978-018-0494-7)