Research Article

Hardcore smoking among daily smokers in male and female adults in 27 countries: a secondary data analysis of Global Adult Tobacco Surveys (2008-2014)

Chandrashekhar T Sreeramareddy¹, Joanne Hon², Anshad Mohamed Abdulla³, Sam Harper⁴

¹ Department of Community Medicine, International Medical University, Kuala Lumpur, Malaysia, ² Clinical School, International Medical University, Seremban, Malaysia, ³ Department of Pediatric Dentistry and Orthodontic Sciences, College of Dentistry, King Khalid University, Abha, Kingdom of Saudi Arabia, ⁴ Department of Epidemiology, Biostatistics and Occupational Health, McGill University, Montreal, Canada

Keywords: global health

<https://doi.org/10.29392/joghr.2.e2018019>

Journal of Global Health Reports

Vol. 2, 2018

Background

“Hardcore smokers” (HCS) who do not want to quit make it more difficult for tobacco control efforts to further reduce smoking prevalence. We aimed to quantify the burden of HCS among daily smoking adult males and females in 27 countries.

Methods

We used Global Adult Tobacco Survey (GATS) data to estimate the prevalence of HCS ie, daily smokers who smoke within 30 minutes after waking up, smoke ≥ 10 cigarettes per day, have not made any quit attempts during previous 12 months or have no intention to quit at all during the coming 12 months. For each GATS country, we estimated sex-wise, weighted and age-adjusted prevalence of daily smoking and HCS.

Results

Overall weighted population prevalence (%) of HCS was highest in Greece (21.0), followed by Russia (13), Poland (9.4), Romania (9.0), and Ukraine (8.9) and lowest in Nigeria (0.4). Estimated number of HCS (in millions) was highest in China (35.8) followed by India (28.2), Russia (18.5), Indonesia (16.1) and lowest in Panama (0.03). The proportion (%) of daily smokers classified as HCS was highest in Greece (56.2%) followed by Russia (42.2%), Ukraine (37.2) and Poland (36.2) and lowest in Mexico (8.29). Overall, proportion of HCS was higher among males in all countries. However, in Greece, Russia, Romania, Ukraine and Poland both population prevalence of HCS among women and proportion of HCS among daily smoking women was higher than in other countries.

Conclusions

At the country-level, a higher daily smoking rates also suggest a higher proportion of HCS. Countries with greater burden of HCS pose greater challenges to tobacco control efforts specifically towards tobacco cessation interventions. Interventions to reduce tobacco use and smoking-related mortality may need to be altered in populations with high proportions of HCS.

Tobacco use, particularly cigarette smoking including environmental tobacco smoke is still a substantial contributor to the global disease burden.¹ Smoking prevalence is decreasing worldwide, particularly in developed countries, although rate of decline has slowed down in recent years.² The sustained presence of heavy smokers who are more addicted and less able to quit³ has led to emergence of “hardening hypotheses”.⁴ Light smokers are more likely to quit than the heavy smokers, thus leading over time to a decrease in light smokers and a relative increase in heavy smokers. Over time these heavy smokers become much harder to reach, and it becomes more difficult for them to quit,⁵ leading to a “hardened” population of smokers who

are more resistant to quit. However, it has also been argued that the process of ‘hardening’ may not occur due to dynamic nature of cohorts of smokers arising from changes in subpopulations of quitters, smokers who die and new smokers.⁶

The ‘hardening hypothesis’ has been tested using data on smoking behaviours in United States,⁷⁻⁹ Australia¹⁰ England,^{11,12} Norway¹³ and Italy.¹⁴ The ‘hardening hypothesis’ has also been evaluated in several studies using varying definitions for ‘hardcore smoker’ (HCS). In general, HCS was defined based on the constructs of duration and intensity of smoking, time to smoke the first cigarette of the day, number of quit attempts made, intention to quit, and

knowledge about harms of smoking.^{6,15} Additionally, empirical evidence for the "hardening hypothesis" has been evaluated using repeated cross-sectional survey data^{16,17} and national population monitoring data.¹⁶ Based on Global Adult Tobacco Survey (GATS), HCS was defined using different constructs and prevalence, and factors associated with HCS were reported.^{18,19} A study from Poland assessed factors associated with HCS including the constructs listed above to define HCS¹⁸ while another study from Bangladesh, India and Thailand tested the association of HCS with socio-demographic factors only.¹⁹

Studying the population prevalence of HCS and the proportion of current daily smokers who are HCS helps design tobacco control policies and set up support services for smoking cessation targeted at the 'hardened' smokers.²⁰ In this paper, we use a five-item construct for HCS based on definitions used in previous studies. We report sex-specific, country-wise proportions of HCS among the current daily smokers, population prevalence of HCS and estimated total number of HCS in 27 GATS countries.

METHODS

We used publicly available data of the Global Adult Tobacco Surveys (GATS) (<http://nccd.cdc.gov/gtssdata/Ancillary/DataReports.aspx?CAID=2>), a series of nationally representative, cross-sectional household surveys done as a part of worldwide Global Tobacco Surveillance System (GTSS) to monitor tobacco use among various population groups. The GATS uses a standardised questionnaire to assess tobacco use behaviours among civilian, non-institutionalised individuals aged 15 years and above.²¹ In each GATS country, the residents of all regions were eligible to be sampled by a stratified multi-stage probability sampling technique. Within a selected geographic location, households were selected at random and within a selected household, all eligible persons were interviewed using a handheld device used for rostering and data collection, and one household member was selected at random for the interview. The implementing agencies in each country adapted the core questionnaire to suit the local tobacco use context. In all countries interviews were done privately by either a male or female interviewers, except for India, Bangladesh, Indonesia and Qatar interviewers were of same gender as the respondent. Further details about survey instrument, methodology etc. are published elsewhere.²¹ The sample sizes, response rates by sex are given in [Table 1](#).

MAIN OUTCOME VARIABLE

Based on previous studies respondents were defined as HCS, if they satisfied the following five criteria⁸: 1) is a current daily smoker; 2) smokes 10 or more cigarettes per day; 3) smokes their first cigarette within 30 minutes after waking up; 4) has not made any quit attempts during 12 months prior to the date surveyed; and 5) has no intention to quit smoking at all or during the next 12 months.

STATISTICAL ANALYSES

All the analyses were done on Stata/IC version 10 (Stata Corp LLC, College Station, TX, USA). Descriptive statistics were calculated for current daily smoking and five constructs of HCS including average of number of cigarettes smoked per day. The proportion of current daily smokers who were HCS was calculated for men, women and both sexes. To adjust for complex sampling design used in GATS, we used country sample weights to estimate prevalence rates and 95% confidence intervals (95% CI). We used total and sex-wise population aged 15 years above from United Nations census data to estimate the number (in millions) male and female HCS in each GATS country. As a sensitivity analysis, we also estimated age-adjusted rates of current daily smoking and HCS, adjusted to the world population standard.²²

RESULTS

[Table 1](#) shows the year of survey followed by sex-specific sample size, median age of respondents and smoking status for each country. A total of 344,329 adults (164,259 men and 180,070 women) were surveyed, median age of the respondents varied between 31-55 years for men and 30-61 years for women. The weighted prevalence of current daily smoking among men was highest in Indonesia (57%) and Russia (55%), and lowest in Panama (4%) and Nigeria (6%). Among women current daily smoking prevalence was considerably lower: <1% in Nigeria, Egypt, Cameroon, Kenya and Malaysia and less than 5% in 12 countries; over 15% in Russia, Uruguay Poland and Greece ([Table 1](#)).

In all countries, the weighted population prevalence HCS was considerably higher among men than women, and in nine countries HCS prevalence was higher than 10% ([Table 2](#)). Among men, Greece (30.9%) had the highest prevalence, followed by Russia (23.8%) while in Argentina, Qatar, Panama, Cameroon, Kenya, Uganda, Mexico and Nigeria HCS prevalence was much lower (2.71-0.81%). However, in 25 countries the weighted population prevalence of HCS among women was less than 5.0% and in 17 of these 25 countries HCS prevalence was <1.0%; Poland (6.8%), and Greece (11.4%) had higher prevalence for women. ([Figure 1](#) shows weighted prevalence of current daily smoking and HCS in each country).

Overall (both sexes), proportion of daily smokers who were defined as HCS was higher in Greece, Cameroon, Russia, Ukraine, Romania and Poland (36.2-56.2%), in other countries the proportion was between 30% and 10%, except Mexico (8.3%). The proportion of male HCS was over 55% in Greece, and in Russia, Ukraine, Poland and Romania it was nearly 40%, whereas in most other countries it ranged between 30-15%. In all countries except Qatar, Bangladesh and Nigeria, the proportion of HCS was higher among males where prevalence of current daily smoking was also low. In both sexes, proportion of daily smokers who were defined as HCS tended to be higher among countries with higher prevalence of daily smoking ([Figure 2](#)).

Overall in 27 GATS countries there were an estimated 111 million HCS (101.6 million men and 12.4 million women, [Table 3](#)). Among the GATS countries, four countries

Table 1. Survey characteristics, sex-wise and country-wise distribution of smoking characteristics and weighted prevalence of daily smoking in 27 GATS countries

		Smoking status (number and %)	Weighted prevalence and 95% CI of daily smoking				
Country	Survey year	Sample size (% of total sample)	Median age (IQR)	Daily smoker	Non-daily smoker	Non-smoker	
Men (≥15 years):							
Argentina	2012	2985 (44.9)	56 (37, 70)	681 (22.8)	271 (9.1)	2033 (68.1)	21.89 (17.32-26.47)
Bangladesh	2009	4468 (46.4)	35 (25, 45)	1972 (44.1)	185 (4.1)	2311 (51.7)	40.69 (38.50-42.87)
Brazil	2008	18039 (45.8)	38 (27, 51)	3513 (19.5)	524 (2.9)	14002 (77.6)	18.91 (18.18-19.63)
Cameroon	2013	4049 (60.9)	31 (23, 42)	79 (2.0)	6 (0.1)	3969 (97.9)	9.11 (7.63-10.61)
China	2010	6603 (49.4)	46 (36, 59)	3303 (50.0)	469 (7.1)	2831 (42.9)	45.35 (42.73-47.97)
Egypt	2009	10062 (48.1)	34 (25, 47)	3904 (38.8)	189 (1.9)	5969 (59.3)	35.85 (34.52-37.17)
Greece	2013	2092 (48.0)	48 (35, 67)	1037 (49.6)	27 (1.3)	1028 (49.1)	49.7 (46.42-52.99)
India	2009	33767 (48.7)	35 (26, 45)	8164 (24.2)	2084 (6.2)	23519 (69.7)	18.35 (17.43-19.27)
Indonesia	2011	3948 (47.5)	38 (28, 52)	2335 (59.1)	385 (9.8)	1228 (31.1)	56.74 (53.82-59.65)
Kazakhstan	2014	2085 (47.1)	39 (28, 52)	795 (38.1)	118 (5.7)	1172 (56.2)	36.91 (34.20-39.63)
Kenya	2014	2077 (47.1)	35 (26, 46)	341 (16.4)	82 (3.9)	1654 (79.6)	11.60 (9.71-13.50)
Malaysia	2011	2104 (49.5)	41 (29, 55)	873 (41.5)	82 (3.9)	1149 (54.6)	39.87 (36.51-43.23)
Mexico	2009	6160 (45.2)	36 (27, 50)	654 (10.6)	744 (12.1)	4762 (77.3)	11.81 (10.61-13.02)
Nigeria	2012	5058 (51.8)	31 (24, 45)	323 (6.4)	91 (1.8)	4644 (91.8)	5.55 (4.60-6.51)
Pakistan	2014	3782 (48.2)	35 (24, 46)	811 (21.4)	67 (1.8)	2904 (76.8)	20.59 (18.88-22.30)
Panama	2013	7679 (45.3)	39 (27, 54)	298 (3.9)	477 (6.2)	6904 (89.9)	4.37 (2.98-5.76)
Philippines	2009	4740 (48.9)	37 (27, 50)	1887 (39.8)	420 (8.9)	2433 (51.3)	38.24 (36.35-40.13)
Poland	2009	3867 (49.3)	47 (32, 60)	1309 (33.9)	116 (3.0)	2442 (63.1)	33.52 (31.58-35.46)
Qatar	2013	4237 (50.5)	33 (26, 41)	799 (18.9)	178 (4.2)	3260 (76.9)	16.51 (14.95-18.07)
Romania	2011	2070 (45.8)	56 (40, 70)	657 (31.7)	50 (2.4)	1363 (65.8)	34.85 (32.27-37.43)
Russia	2009	6217 (54.5)	46 (32, 58)	3486 (56.1)	300 (4.8)	2431 (39.1)	54.98 (53.15-56.80)
Thailand	2011	8781 (42.6)	47 (35, 60)	3525 (40.1)	368 (4.2)	4888 (55.7)	42.04 (40.17-43.90)
Turkey	2012	4470 (45.4)	43 (30, 59)	1593 (35.6)	189 (4.2)	2688 (60.1)	37.33 (35.38-39.27)
Uganda	2013	3853 (45.3)	31 (24, 42)	400 (10.4)	79 (2.1)	3374 (87.6)	8.69 (7.49-9.88)

	Smoking status (number and %)	Weighted prevalence and 95% CI of daily smoking					
Ukraine	2010	4076 (50.0)	51 (35, 66)	1866 (45.8)	168 (4.1)	2042 (50.1)	45.48 (43.55-47.420)
Uruguay	2009	2634 (47.2)	47 (32, 65)	677 (25.7)	134 (5.1)	1823 (69.2)	24.85 (22.45-27.25)
Viet Nam	2010	4356 (43.9)	40 (29,53)	1794 (41.2)	377 (8.7)	2185 (50.2)	38.74 (36.88-40.61)
Women (≥15 years):							
Argentina	2012	3660 (55.1)	37 (27, 54)	505 (13.8)	193 (5.3)	2962 (80.9)	12.74 (9.79-15.69)
Bangladesh	2009	5161 (53.6)	37 (27, 48)	66 (1.3)	10 (0.2)	5085 (98.5)	1.33 (0.87-1.79)
Brazil	2008	21386 (54.2)	38 (27,54)	2560 (12.0)	406 (1.9)	18420 (86.1)	11.52 (10.98-12.06)
Cameroon	2013	2594 (39.1)	30 (23, 44)	18 (0.7)	8 (0.3)	2568 (99.0)	0.46 (0.20-0.71)
China	2010	6751 (50.6)	47 (37, 60)	197 (2.9)	41 (0.6)	6513 (96.5)	2.0 (1.44-2.55)
Egypt	2009	10862 (51.9)	38 (28, 49)	53 (0.5)	9 (0.1)	10800 (99.4)	0.45 (0.24-0.67)
Greece	2013	2267 (52)	47 (34, 65)	573 (25.3)	30 (1.3)	1664 (73.4)	23.91 (21.05-26.76)
India	2009	35529 (51.3)	36 (27, 48)	1059 (3.0)	289 (0.8)	34181 (96.2)	2.44 (2.06-2.82)
Indonesia	2011	4357 (52.5)	38 (29, 50)	90 (2.1)	45 (1.0)	4222 (96.9)	1.82 (1.29-2.34)
Kazakhstan	2014	2340 (52.9)	41 (29, 56)	77 (3.3)	30 (1.3)	2233 (95.4)	3.17 (2.24-4.10)
Kenya	2014	2331 (52.9)	31 (24, 45)	20 (0.9)	6 (0.3)	2305 (98.9)	0.60 (0.14-1.06)
Malaysia	2011	2146 (50.5)	40 (28, 53)	25 (1.2)	9 (0.4)	2112 (98.4)	0.69 (0.32-1.06)
Mexico	2009	7457 (54.8)	37 (26, 52)	202 (2.7)	221 (3.0)	7034 (94.3)	3.67 (2.88-4.47)
Nigeria	2012	4707 (48.2)	34 (25, 45)	12 (0.3)	3 (0.1)	4692 (99.7)	0.28 (0.099-0.46)
Pakistan	2014	4049 (51.8)	32 (24, 44)	79 (2.0)	6 (0.1)	3964 (97.9)	1.99 (0.02-2.52)
Panama	2013	9283 (54.7)	42 (28, 58)	69 (0.7)	152 (1.6)	9062 (97.6)	1.24 (0.69-1.79)
Philippines	2009	4961 (51.1)	36 (27, 48)	360 (7.3)	102 (2.1)	4499 (90.7)	6.86 (5.94-7.78)
Poland	2009	3973 (50.7)	44 (31, 57)	870 (21.9)	121 (3.0)	2982 (75.1)	21.0 (19.54-22.47)
Qatar	2013	4161 (49.5)	38 (29, 45)	52 (1.2)	50 (1.2)	4059 (97.5)	1.66 (1.02-2.300)
Romania	2011	2447 (54.2)	50 (36, 65)	301 (12.3)	48 (2.0)	2098 (85.7)	14.49 (12.70-16.29)
Russia	2009	5189 (45.5)	43 (29, 55)	786 (15.1)	235 (4.5)	4168 (80.3)	16.26 (14.38-18.14)
Thailand	2011	11825 (57.4)	46 (33, 58)	328 (2.8)	70 (0.6)	11427 (96.6)	2.11 (1.75-2.47)
Turkey	2012	5381 (54.6)	42 (31, 56)	520 (9.7)	110 (2.0)	4751 (88.3)	10.67 (9.59-11.75)

	Smoking status (number and %)	Weighted prevalence and 95% CI of daily smoking					
Uganda	2013	4655 (54.7)	30 (23, 43)	71 (1.5)	19 (0.4)	4565 (98.1)	1.31 (0.91-1.7)
Ukraine	2010	4082 (50.0)	46 (32, 59)	289 (7.1)	74 (1.8)	3719 (91.1)	8.88 (7.60-10.15)
Uruguay	2009	2947 (52.8)	45 (31, 61)	488 (16.6)	95 (3.2)	2364 (80.2)	16.42 (14.77-18.07)
Viet Nam	2010	5569 (56.1)	40 (28, 52)	73 (1.3)	16 (0.3)	5480 (98.4)	1.21 (0.75-1.68)

CI – confidence interval, IQR – interquartile range

Figure 1

Proportions of hardcore smokers (HCS) among daily smoking men and women in 27 GATS countries.

together accounted for nearly 70% of all HCS: China (30 million), India (21 million), Russia (16 million) and Indonesia (11 million). China and India had largest burden of HCS in terms of absolute number of HCS due to their large population sizes. India and China had relatively lower prevalence of current daily smoking than Greece and eastern European countries of GATS which had very high current daily smoking prevalence. Age-standardization led to similar estimated prevalence for both daily smoking and HCS, compared to crude estimates (see Table S1 in **Online Supplementary Document(Online Supplementary Document)**).

DISCUSSION

The prevalence of HCS and the proportion of daily smokers defined as ‘hardcore’ varied widely between 27 GATS coun-

Figure 2

Scatter diagram for relationship of percentage of hardcore smokers (HCS) and prevalence of daily smoker among men and women in 27 GATS countries. Comparison of prevalence of daily smoker with percentage of HCS by each country can also be found in Table S1 in **Online Supplementary Document**.

tries and sexes. In general, we found that proportion of daily smokers defined as “hardcore” was higher among countries with a higher prevalence of current daily smoking. In terms of absolute numbers China, India, Russia and Indonesia have very large numbers of HCS, whereas Russia has highest proportion of daily smokers defined as “hardcore”. In all GATS countries, women ranked lower than men in prevalence of HCS and number of HCS. Both prevalence of HCS and the proportion of HCS among current daily smokers were higher in high-income and upper-middle-income GATS countries, Qatar being an exception to this.

A strength of our analysis was a robust survey design of GATS, which generates nationally representative samples of men and women. The standardised methodology and survey instrument of GATS enabled us to construct a definition of HCS for cross-country comparison.²³ Nevertheless, es-

Table 2. Weighted prevalence rates of HCS and distribution (number and %) of the HCS constructs among the daily smokers in 27 GATS countries

	Weighted prevalence of HCS (95% CI)	Mean sticks per day	≥10 sticks per day	No quit attempt in last 12 months	No intention to quit at all or next 12 months	Smoke within 30 minutes after waking up
Men (≥15 years)						
Argentina	2.31 (1.54-3.08)	15.26	499 (73.3)	554 (58.2)	721 (75.7)	219 (32.2)
Bangladesh	7.46 (6.21-8.71)	12.33	1220 (61.9)	1126 (52.2)	1358 (63)	881 (44.7)
Brazil	4.67 (4.29-5.05)	14.96	2316 (68.1)	2546 (60.8)	1977 (80.5)	2195 (62.5)
Cameroon	1.55 (0.86, 2.23)	8.76	95 9 (33.9)	218 (58.0)	258 (68.6)	83 (28.9)
China	5.12 (4.40-5.85)	17.24	2694 (81.6)	1044 (69.6)	3172 (84.1)	1840 (55.7)
Egypt	6.4 (5.72-7.08)	16.74	3114 (79.8)	2541 (62.1)	3004 (73.4)	1402 (35.9)
Greece	30.91 (27.78-34.04)	20	866 (83.5)	906 (85.2)	927 (87.1)	795 (76.7)
India	4.46 (4.03-4.90)	10.62	3721 (45.6)	7159 (69.9)	8004 (78.1)	5370 (65.8)
Indonesia	13.09 (10.95-15.24)	12.38	1545 (66.2)	1948 (71.6)	2427 (89.2)	906 (38.8)
Kazakhstan	10.52 (8.83, 12.21)	16.73	552 (77.0)	675 (73.9)	775 (84.9)	422 (53.1)
Kenya	1.97 (1.07, 2.86)	9.36	142 (42.8)	193 (45.6)	241 (57.0)	256 (75.1)
Malaysia	8.15 (6.46-9.85)	14.38	654 (74.9)	564 (59.1)	842 (88.2)	417 (47.8)
Mexico	1.15 (0.78-1.52)	10.02	268 (41.0)	789 (56.4)	922 (66.0)	180 (27.5)
Nigeria	0.81 (0.38-1.23)	8.9	105 (32.5)	235 (56.8)	284 (68.6)	168 (52.0)
Pakistan	5.54 (4.60, 6.48)	13.69	521 (65.4)	668 (76.1)	740 (84.3)	399 (49.2)
Panama	1.53 (0.83-2.23)	12.77	118 (39.6)	393 (50.7)	600 (77.4)	130 (43.6)
Philippines	10.49 (9.26-11.73)	11.48	1176 (62.3)	1259 (54.6)	1831 (79.4)	1124 (59.6)
Poland	12.27 (11.04-13.51)	17.84	1141 (87.2)	1006 (70.6)	992 (69.6)	817 (62.4)
Qatar	2.71 (2.08-3.33)	14.64	554 (69.3)	635 (65.0)	601 (61.5)	298 (37.3)
Romania	13.74 (11.87-15.61)	17.04	567 (86.3)	466 (65.9)	547 (77.4)	471 (71.7)
Russia	23.8 (22.01-25.58)	17.77	2892 (83.0)	2727 (72.0)	3334 (88.1)	2357 (67.6)
Thailand	12.19 (10.98-13.39)	12.24	2267 (64.3)	2535 (65.1)	3303 (84.8)	2093 (59.4)
Turkey	8.05 (7.00-9.11)	20.36	1398 (87.8)	1034 (58.0)	1143 (64.1)	666 (41.8)
Uganda	0.98 (0.59, 1.38)	6.95	287 (59.9)	192 (48.0)	346 (72.2)	278 (69.5)
Ukraine	16.65 (15.25-18.06)	17.25	1622 (86.9)	1343 (66.0)	1552 (76.3)	1231 (66.0)
Uruguay	4.37 (3.24-5.50)	17.11	500	465 (57.3)	550 (67.7)	285 (42.1)

	Weighted prevalence of HCS (95% CI)	Mean sticks per day	≥10 sticks per day	No quit attempt in last 12 months	No intention to quit at all or next 12 months	Smoke within 30 minutes after waking up
			(73.9)			
Viet Nam	9.88 (8.84-10.92)	14.92	1268 (70.8)	1037 (47.8)	1550 (71.4)	1167 (65.1)
Women (≥15 years):						
Argentina	1.59 (0.78-2.39)	11.39	296 (58.6)	378 (54.2)	515 (73.8)	145 (28.7)
Bangladesh	3.82 (3.16-4.49)	8.57	21 (31.8)	49 (64.5)	60 (78.9)	33 (50.0)
Brazil	2.33 (2.08-2.59)	12.39	1474 (59.6)	1619 (54.6)	1272 (78.6)	1399 (54.6)
Cameroon	0.09 (-0.001, 0.19)	5.35	4 (23.5)	9 (34.6)	19 (73.1)	9 (50.0)
China	0.12 (0.04-0.19)	12.64	135 (68.5)	49 (59.8)	196 (82.4)	105 (53.3)
Egypt	0.05 (0.0027-0.09)	7.98	17 (32.1)	34 (54.8)	46 (74.2)	22 (41.5)
Greece	11.39 (9.57-13.21)	15	471 (82.2)	501 (83.1)	516 (85.6)	367 (64.0)
India	0.31 (0.21-0.40)	7.27	275 (26.0)	939 (69.7)	1086 (80.6)	625 (59.0)
Indonesia	0.22 (0.0591-0.37)	8.01	29 (32.2)	86 (63.7)	121 (89.6)	25 (27.8)
Kazakhstan	0.38 (0.06, 0.7)	10.49	30 (49.2)	75 (70.1)	85 (79.4)	34 (44.2)
Kenya	0.013 (-0.01, 0.04)	5.8	2 (25.0)	9 (34.6)	10 (38.5)	14 (70.0)
Malaysia	0.03 (0.012-0.07)	9.84	15 (60)	19 (55.9)	28 (82.4)	7 (28)
Mexico	0.22 (0.051-0.38)	8.35	56 (27.7)	192 (45.4)	252 (59.6)	62 (30.7)
Nigeria	0.042 (0.017-0.10)	18.83	5 (41.7)	8 (53.3)	12 (80.0)	7 (58.3)
Pakistan	0.2 (0.02, 2.5)	7.45	23 (30.3)	63 (74.1)	74 (87.1)	23 (29.1)
Panama	0.2 (0.047-0.36)	7.8	19 (27.5)	112 (50.7)	183 (82.8)	25 (36.2)
Philippines	0.74 (0.44-1.03)	6.88	116 (32.2)	222 (48.1)	349 (75.5)	163 (45.3)
Poland	6.78 (5.91-7.66)	15.49	742 (85.3)	676 (68.2)	654 (66.0)	502 (57.7)
Qatar	0.64 (0.20-1.08)	11.65	26 (50.0)	60 (58.8)	69 (67.6)	26 (50.0)
Romania	4.6 (3.47-5.72)	13.52	226 (75.1)	215 (61.6)	263 (75.4)	188 (62.5)
Russia	3.99 (3.14-4.85)	11.43	532 (67.7)	677 (66.3)	855 (83.7)	369 (46.9)
Thailand	0.45 (0.30-0.60)	8.61	140 (42.7)	253 (63.6)	320 (80.4)	163 (49.7)
Turkey	1.98 (1.53-2.42)	15.12	361 (69.4)	350 (55.6)	382 (60.6)	200 (38.5)
Uganda	0.06 (-0.013, 0.14)	3.47	4 (5.6)	52 (57.8)	71 (78.9)	46 (64.8)
Ukraine	2.39 (1.69-3.08)	11.3	189	222 (61.2)	260 (71.6)	150 (51.9)

	Weighted prevalence of HCS (95% CI)	Mean sticks per day	≥10 sticks per day	No quit attempt in last 12 months	No intention to quit at all or next 12 months	Smoke within 30 minutes after waking up
			(65.4)			
Uruguay	2.3 (1.53-3.06)	12.96	312 (64.2)	317 (54.4)	384 (65.9)	158 (32.4)
Viet Nam	0.31 (0.14-0.49)	10.25	30 (41.1)	55 (61.8)	67 (75.3)	40 (54.8)

HCS – hardcore smoking, CI – confidence interval

Table 3. Estimated number of HCS and their 95CIs in 27 GATS countries by both sexes and sex-wise

Country	Total number HCS (both sexes)			Number of male HCS			Number of female HCS
	Estimates	95% CIs		Estimates	95% CIs		
Argentina	600,767	426,451	775,083	347,841	231,894	463,788	255,510
Bangladesh	3,596,984	3,154,423	4,048,961	3,471,427	2,889,753	4,053,100	1,819,390
Brazil	3,388,870	3,012,329	3,750,929	3,270,889	3,004,735	3,537,043	1,742,445
Cameroon	102,667	59,033	145,017	99,426	55,165	143,045	5,777
China	29,676,736	25,786,527	33,566,945	28,801,771	24,751,522	32,908,274	658,744
Egypt	1,630,698	1,481,549	1,779,847	1,620,481	1,448,305	1,792,657	11,222
Greece	1,980,614	1,821,070	2,140,158	1,416,617	1,273,168	1,560,066	553,263
India	20,541,066	18,696,562	22,469,411	19,129,782	17,285,431	21,017,025	1,269,425
Indonesia	11,224,127	10,243,707	12,221,452	11,039,063	9,234,358	12,852,202	186,353
Kazakhstan	393,981	328,093	461,213	351,459	291,825	411,093	14,205
Kenya	252,969	138,220	365,110	255,229	138,627	370,535	1,706
Malaysia	863,312	705,414	1,021,210	859,365	681,165	1,038,619	2,989
Mexico	526,817	383,140	678,476	441,100	299,181	583,019	91,221
Nigeria	351,291	171,561	522,851	331,155	155,357	502,865	17,141
Pakistan	3,831,873	3,191,048	4,485,777	3,720,310	3,089,066	4,351,554	17,039
Panama	20,982	11,817	29,906	18,435	10,001	26,869	2,462
Philippines	3,449,610	3,092,329	3,806,891	3,227,957	2,849,464	3,609,527	228,130
Poland	3,039,755	2,803,689	3,279,055	1,894,431	1,704,525	2,085,882	1,145,704
Qatar	29,551	23,401	35,358	35,948	27,591	44,173	2,442
Romania	1,523,880	1,341,014	1,708,438	1,119,847	967,437	1,272,257	403,960
Russia	15,649,954	14,684,652	16,603,190	13,031,482	12,051,383	14,006,105	2,629,798
Thailand	3,277,696	3,011,216	3,544,175	3,150,351	2,837,642	3,460,475	123,535
Turkey	2,798,306	2,477,374	3,113,608	2,258,669	1,964,060	2,556,084	559,270
Uganda	88,860	53,316	124,403	86,213	51,904	121,403	5,385
Ukraine	3,485,824	3,195,011	3,776,637	2,959,780	2,710,910	3,210,428	514,390
Uruguay	84,635	68,895	100,376	53,530	39,688	67,372	31,174
Viet Nam	3,216,745	2,879,506	3,553,985	3,118,793	2,790,499	3,447,086	103,190
Total	115,628,570	103,241,349	128,108,462	10,6111,349	92,834,654	119,492,544	12,395,870

HCS – hardcore smoking, CI – confidence interval

imates reported in this paper should be interpreted with caution against the possible limitations inherent in GATS design. Self-reported smoking behaviour is known to lead to under-estimates due to smoking-related stigma.²⁴ Furthermore, social desirability bias related to quit attempts and intentions to quit may have resulted in underestimation of HCS.²⁵ Both duration of smoking as well as smoking intensity (ie, daily number of cigarettes smoked) indicate nico-

tine addiction but duration of smoking could not be computed in many low-income and lower-middle-income GATS countries because of missing data on age of smoking initiation. The surveys we analysed were implemented at different time periods, smoking patterns change over time and are sensitive to country-specific tobacco control policies²; hence our results should be interpreted in the context of the local tobacco control environment in the GATS countries.

The prevalence of HCS among high-income countries has been well reported and ranged from 16% in the UK¹¹ to 6% in Norway.¹³ Our estimates varied widely and ranged from 21% in Greece to 0.4% in Nigeria. However, our estimates of HCS are not comparable to those from high-income countries due to heterogeneity in the constructs these studies used to define a HCS.¹⁵ Given the relationship, we identified between daily rates of smoking and HCS, our estimates may also differ from high-income countries because the prevalence of daily smoking differs as well. Two prior GATS-based studies used different definitions for HCS and reported a population prevalence of HCS of 3.1% (India), 3.8% (Bangladesh) and 6.0% (Thailand)¹⁹ whereas 10.0% in Poland¹⁸ which were nearly same as our estimates.

The hardening hypothesis has implications for tobacco control, since when the prevalence of smoking decreases it would make sense to target HCS to achieve further reductions.⁵ Studies on HCS have primarily been based on cross-sectional data.^{12,14,18,19} Additional studies that have examined sequential cross-sectional surveys have provided some evidence against the ‘hardening theory’^{13,17,26}; however, one study reported that hardening may be occurring in the UK.⁵ Nevertheless, we are only able to provide cross-sectional estimates of HCS for 27 GATS countries. Some GATS countries have a huge burden of estimated population who are HCS, mainly among men in China, India, Russia and Indonesia. These estimate numbers have implications for tobacco control since smoking cessation can help to avoid a substantial tobacco-attributable mortality, particularly in low-and-middle-income countries (LMIC) currently experiencing epidemic levels of smoking.²⁷ Moreover, these numbers have implications for healthcare services in LMICs which typically neither provided smoking cessation services²⁸ nor implemented tobacco-related training programs healthcare professionals²⁹ as per the recommendations of FCTC Article 14 or the FCTC Article 14 guidelines. LMICs with higher burden of HCS and low quit rates are likely to require larger healthcare costs to implement smoking cessation services.

Tobacco control strategies should consider smoking intensity, duration of smoking, time to smoke first cigarette, as these factors are known to be associated with tobacco dependence.³⁰ More intensive and universal measures should be undertaken to achieve higher quit rates among all daily smokers.⁵ Reaching out to the hardcore smokers who lack motivation (don’t want to quit) and have severe nicotine addiction (can’t quit)¹⁷ requires a more holistic approach. Motivational interviewing maybe one of the approaches to help smokers decide to quit,³¹ as it is well known that smokers who are in contemplation stages are more likely to remain tobacco-free.³² HCS constructs in most studies including ours do not include socio-economic deprivation and mental health which are associated with “hardening”.¹⁴ As smoking is also known to be common among those with socio-economic deprivation³³ wider community level interventions, such as health promotion and socio-economic development may not only help cessation but could also potentially help to decrease smoking initiation.³⁴

In addition to conflicting reports on the ‘hardening hypotheses’, there is also a lack of consensus on uniform definitions of hardening across the published literature. Hence,

there is a need for further studies defining HCS based on all the four domains of constructs described by Edwards et al.¹⁷ Current conflicting literature, mainly from developed countries, has tested the ‘hardening hypothesis’ with sequential cross-sectional surveys^{10,11,13,16,17,26}; cohort studies on daily smokers who cannot quit would confirm the hardening hypothesis and examine the changes in proportion of HCS over time.

CONCLUSIONS

HCS constitute a fifth to third of daily smokers in some GATS countries, which presents challenges to tobacco control efforts. As the proportion of HCS tends to increase with the prevalence of daily smoking, efforts should be made to counter smoking initiation. Further HCS also poses a challenge to health services in LMIC where smoking cessation services are sub optimal. Future studies should include all four domains of HCS to better understand about “hardening hypothesis” in LMIC.

ACKNOWLEDGEMENTS

The authors thank the Centre for Disease Control for making the datasets of Global Adult Tobacco Survey that enables us prepare this manuscript.

ETHICS APPROVAL

All the Global Adult Tobacco Surveys were approved by ethical boards of surveys countries and CDC, Atlanta. The data used for this manuscript are available in the public domain are de-identified. Therefore, a separate ethical approval was not required for this manuscript preparation.

FUNDING

This study did not receive any funding.

COMPETING INTERESTS

The authors have completed the Unified Competing Interest form at http://www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare no competing interests.

CORRESPONDENCE TO:

Chandrashekhar T Sreeramareddy
Department of Community Medicine
International Medical University
126, Jalan Jalil Perkasa 19
Bukit Jalil
57000 Kuala Lumpur
Malaysia
chandrashekharats@yahoo.com

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-4.0). View this license's legal deed at <http://creativecommons.org/licenses/by/4.0> and legal code at <http://creativecommons.org/licenses/by/4.0/legalcode> for more information.

REFERENCES

1. Gakidou E, Afshin A, Abajobir AA, et al. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2016: A systematic analysis for the Global Burden of Disease Study 2016. *Lancet*. 2017;390(10100):1345–1422. doi:[10.1016/s0140-6736\(17\)32366-8](https://doi.org/10.1016/s0140-6736(17)32366-8)
2. Reitsma MB, Fullman N, Ng M, et al. Smoking prevalence and attributable disease burden in 195 countries and territories, 1990–2015: A systematic analysis from the Global Burden of Disease Study 2015. *Lancet*. 2017;389(10082):1885–1906. doi:[10.1016/s0140-6736\(17\)30819-x](https://doi.org/10.1016/s0140-6736(17)30819-x)
3. DiFranza JR, Guerrera MP. Hard-core smokers. *JAMA*. 1989;261(18):2634. doi:[10.1001/jama.1989.03420180058016](https://doi.org/10.1001/jama.1989.03420180058016)
4. Darville A, Hahn EJ. Hardcore smokers: what do we know? *Addict Behav*. 2014;39(12):1706–1712. doi:[10.1016/j.addbeh.2014.07.020](https://doi.org/10.1016/j.addbeh.2014.07.020)
5. Docherty G, McNeill A. The hardening hypothesis: does it matter? *Tob Control*. 2012;21(2):267–268. doi:[10.1136/tobaccocontrol-2011-050382](https://doi.org/10.1136/tobaccocontrol-2011-050382)
6. Hughes JR. The hardening hypothesis: is the ability to quit decreasing due to increasing nicotine dependence? A review and commentary. *Drug Alcohol Depend*. 2011;117(2-3):111–117. doi:[10.1016/j.drugalcdep.2011.02.009](https://doi.org/10.1016/j.drugalcdep.2011.02.009)
7. Augustson E, Marcus S. Use of the current population survey to characterize subpopulations of continued smokers: a national perspective on the “hardcore” smoker phenomenon. *Nicotine Tob Res*. 2004;6(4):621–629. doi:[10.1080/14622200410001727876](https://doi.org/10.1080/14622200410001727876)
8. Emery S, Gilpin EA, Ake C, Farkas AJ, Pierce JP. Characterizing and identifying “hard-core” smokers: Implications for further reducing smoking prevalence. *Am J Public Health*. 2000;90(3):387–394. doi:[10.2105/ajph.90.3.387](https://doi.org/10.2105/ajph.90.3.387)
9. Sorg A, Xu J, Doppalapudi SB, Shelton S, Harris JK. Hardcore smokers in a challenging tobacco control environment: the case of Missouri. *Tob Control*. 2011;20(5):388–390. doi:[10.1136/tc.2010.039743](https://doi.org/10.1136/tc.2010.039743)
10. Mathews R, Hall WD, Gartner CE. Is there evidence of ‘hardening’ among Australian smokers between 1997 and 2007? Analyses of the Australian National Surveys of Mental Health and Well-Being. *Aust N Z J Psychiatry*. 2010;44(12):1132–1136. doi:[10.1097/00048674.2010.520116](https://doi.org/10.1097/00048674.2010.520116)
11. Docherty G, McNeill A, Gartner C, Szatkowski L. Did hardening occur among smokers in England from 2000 to 2010? *Addiction*. 2013;109(1):147–154. doi:[10.1111/add.12359](https://doi.org/10.1111/add.12359)
12. Jarvis MJ, Wardle J, Waller J, Owen L. Prevalence of hardcore smoking in England, and associated attitudes and beliefs: cross sectional study. *BMJ*. 2003;326(7398):1061–0. doi:[10.1136/bmj.326.7398.1061](https://doi.org/10.1136/bmj.326.7398.1061)
13. Lund M, Lund KE, Kvaavik E. Hardcore smokers in Norway 1996–2009. *Nicotine Tob Res*. 2011;13(11):1132–1139. doi:[10.1093/ntr/ntr166](https://doi.org/10.1093/ntr/ntr166)
14. Ferketich AK, Gallus S, Colombo P, Pacifici R, Zuccaro P, La Vecchia C. Hardcore smoking among Italian men and women. *Eur J Cancer Prev*. 2009;18(2):100–105. doi:[10.1097/cej.0b013e32830c8d98](https://doi.org/10.1097/cej.0b013e32830c8d98)
15. Costa ML, Cohen JE, Chaiton MO, Ip D, McDonald P, Ferrence R. “Hardcore” definitions and their application to a population-based sample of smokers. *Nicotine Tob Res*. 2010;12(8):860–864. doi:[10.1093/ntr/ntq103](https://doi.org/10.1093/ntr/ntq103)
16. Bommel J, Nagelhout GE, Kleinjan M, Schoenmakers TM, Willemsen MC, van de Mheen D. Prevalence of hardcore smoking in the Netherlands between 2001 and 2012: a test of the hardening hypothesis. *BMC Public Health*. 2016;16(1):754. doi:[10.1186/s12889-016-3434-x](https://doi.org/10.1186/s12889-016-3434-x)
17. Edwards R, Tu D, Newcombe R, Holland K, Walton D. Achieving the tobacco endgame: evidence on the hardening hypothesis from repeated cross-sectional studies in New Zealand 2008–2014. *Tob Control*. 2017;26(4):399–405. doi:[10.1136/tobaccocontrol-2015-052860](https://doi.org/10.1136/tobaccocontrol-2015-052860)
18. Kaleta D, Usidame B, Dziankowska-Zaborszczyk E, Makowiec-Dbrowska T, Leinsalu M. Prevalence and factors associated with hardcore smoking in Poland: findings from the Global Adult Tobacco Survey (2009–2010). *BMC Public Health*. 2014;14(1):583. doi:[10.1186/1471-2458-14-583](https://doi.org/10.1186/1471-2458-14-583)
19. Kishore J, Jena PK, Bandyopadhyay C, Swain M, Das S, Banerjee I. Hardcore smoking in three South-East asian countries: results from the global adult tobacco survey. *Asian Pac J Cancer Prev*. 2013;14(2):625–630. doi:[10.7314/apjcp.2013.14.2.625](https://doi.org/10.7314/apjcp.2013.14.2.625)

20. Warner K, Burns D. Hardening and the hard-core smoker: concepts, evidence, and implications. *Nicotine Tob Res.* 2003;5(1):37-48. [doi:10.1080/1462220021000060428](https://doi.org/10.1080/1462220021000060428)
21. World Health Organization. *Global Adult Tobacco Survey*. World Health Organization; 2009.
22. Ahmadi B, Boschi-Pinto C, Lopez A, Murray C, Lozano R, Inoue M. *Age Standardization of Rates: A New WHO Standard*. World Health Organisation; 2001.
23. Kalsbeek WD, Bowling JM, Hsia J, Mirza S, Palipudi KM, Asma S. The Global Adult Tobacco Survey (GATS): sample design and related methods. Published online 2010.
24. Connor Gorber S, Schofield-Hurwitz S, Hardt J, Levasseur G, Tremblay M. The accuracy of self-reported smoking: a systematic review of the relationship between self-reported and cotinine-assessed smoking status. *Nicotine Tob Res.* 2009;11(1):12-24. [doi:10.1093/ntr/ntn010](https://doi.org/10.1093/ntr/ntn010)
25. Etter JF, Bergman MM, Humair JP, Perneger TV. Development and validation of a scale measuring self-efficacy of current and former smokers. *Addiction.* 2000;95(6):901-913. [doi:10.1046/j.1360-0443.2000.9569017.x](https://doi.org/10.1046/j.1360-0443.2000.9569017.x)
26. Kulik MC, Glantz SA. The smoking population in the USA and EU is softening not hardening. *Tob Control.* 2016;25(4):470-475. [doi:10.1136/tobaccocontrol-2015-052329](https://doi.org/10.1136/tobaccocontrol-2015-052329)
27. Jha P, Peto R. Global effects of smoking, of quitting, and of taxing tobacco. *N Engl J Med.* 2014;370(1):60-68. [doi:10.1056/nejmra1308383](https://doi.org/10.1056/nejmra1308383)
28. Piné-Abata H, McNeill A, Murray R, Bitton A, Rigotti N, Raw M. A survey of tobacco dependence treatment services in 121 countries. *Addiction.* 2013;108(8):1476-1484. [doi:10.1111/add.12172](https://doi.org/10.1111/add.12172)
29. Kruse GR, Rigotti NA, Raw M, et al. Tobacco dependence treatment training programs: an international survey. *Nicotine Tob Res.* 2015;18(5):1012-1018. [doi:10.1093/ntr/ntv142](https://doi.org/10.1093/ntr/ntv142)
30. Fagerström K, Furberg H. A comparison of the Fagerström Test for Nicotine Dependence and smoking prevalence across countries. *Addiction.* 2008;103(5):841-845. [doi:10.1111/j.1360-0443.2008.02190.x](https://doi.org/10.1111/j.1360-0443.2008.02190.x)
31. Heckman CJ, Eggleston BL, Hofmann MT. Efficacy of motivational interviewing for smoking cessation: a systematic review and meta-analysis. *Tob Control.* 2010;19(5):410-416. [doi:10.1136/tc.2009.033175](https://doi.org/10.1136/tc.2009.033175)
32. DiClemente CC, Prochaska JO, Fairhurst SK, Velicer WF, Velasquez MM, Rossi JS. The process of smoking cessation: an analysis of precontemplation, contemplation, and preparation stages of change. *J Consult Clin Psychol.* 1991;59(2):295-304. [doi:10.1037/0022-006x.59.2.295](https://doi.org/10.1037/0022-006x.59.2.295)
33. Sreeramareddy CT, Harper S, Ernsten L. Educational and wealth inequalities in tobacco use among men and women in 54 low-income and middle-income countries. *Tob Control.* 2018;27(1):26-34. [doi:10.1136/tobaccocontrol-2016-053266](https://doi.org/10.1136/tobaccocontrol-2016-053266)
34. Cummings KM. Community-wide interventions for tobacco control. *Nicotine Tob Res.* 1999;1(1):113-116. [doi:10.1080/14622299050011701](https://doi.org/10.1080/14622299050011701)

SUPPLEMENTARY MATERIALS

Online Supplementary Document

Download: <https://www.joghr.org/article/11926-hardcore-smoking-among-daily-smokers-in-male-and-female-adults-in-27-countries-a-secondary-data-analysis-of-global-adult-tobacco-surveys-2008-2014/attachment/29407.pdf>